

DOI: <https://doi.org/10.20372/afnr.v1i2.668>

ISSN: 2520-7687 (Print) and 3005-7515 (Online)

Journal of Agriculture, Food and Natural Resources

J. Agric. Food Nat. Resour., Oct - Dec 2023,1(2):01-11

Journal Homepage: <https://journals.wgu.edu.et>

Original Research

**Minimum Acceptable Diet and Associated Factors Among Infants and Young Children Aged 6-23 Months in Jigjiga, Ethiopia**Abdirauf Ahmed Isse<sup>1</sup>, Zelalem Tafese<sup>2</sup>, Abdulkarim Mohamed<sup>1</sup>, Shamsedin Mahdi Hassan<sup>1\*</sup>, and Ziad Ahmed<sup>1</sup><sup>1</sup>College of Dryland Agriculture, Department of Food Science and Nutrition, Jigjiga University, Ethiopia<sup>2</sup>School of Nutrition, Food Science and Technology, Hawassa University, Ethiopia**Abstract****Article Information**

A minimum acceptable diet (MAD) is one of the valid and reliable indicators to assess complementary feeding practices in 6–23-month-old children. The study aimed to assess the proportion of MAD and associated factors among young children aged 6–23 months in Jigjiga City, Ethiopia, 2021. A sample of 292 mothers with children aged 6–23 months were randomly selected from six kebeles. A systematic random sampling technique was used to identify study participants. Data were collected using a pre-tested structured questionnaire. A logistic regression model was applied to identify the significant factors associated with MAD. The result showed that 22.2% of the children between the ages of six and twenty-three months met the required MAD. The place of delivery [AOR=1.99, 95% CI (1.07-3.70)], aged group of 12–17 months [AOR=2.61, 95% CI (1.15-5.92)], aged group of 18-23 months [AOR=3.2 95% CI (1.25-8.17)] and receiving child feeding counseling [AOR=2.54, 95% CI (1.21-5.35)] were significantly associated with adequate MAD compared to their counterparts. The study found that the proportion of children aged 6-23 months who met the recommended MAD was low in Jigjiga City, Ethiopia.

**Article History:****Received:** 14-08-2023**Revised:** 03-12-2023**Accepted:** 08-12-2023**Keywords:**Child feeding,  
Ethiopia,  
Factors,  
Jigjiga,  
Minimum acceptable diet**\*Corresponding Author:****E-mail:** shamsedinmahdi1@gmail.com

Copyright©2023 AFNR Journal, Wallaga University. All Rights Reserved

**INTRODUCTION**

The first 1000 days, or the period from conception to the end of a child's second year, have come to be known as the earliest stage of human development. Governments, the early years sector, and some commercial organizations have increased their attention on the first 1000 days as a result of the importance of this time frame for long-term health and developmental outcomes (Moore *et al.*, 2017; Prado & Dewey, 2014). Consequently, proper nutrition at the early age of the child is the cornerstone for developing the maximum human potential in adult life WHO (2018). For this, a child should fulfill the timely introduction of complementary feeding, and meet the minimum acceptable diet (MAD) adequate for his age WHO (2010; 2018). Since MAD is a composite indicator of minimum dietary diversity (MDD) and minimum meal frequency (MMF) WHO (2010). MDD is utilized to assess the quality of a baby's and young child's complementary diet while MMF is used as a stand-in measure of caloric intake or quantity of food consumed other than breast milk. Furthermore, MAD assesses both micronutrient adequacy and

quantity of food consumed during the previous day or night and measures appropriate complementary feeding practices WHO (2010). Thus, the MAD assesses several aspects of the diet of newborns and young children. Optimal nutrition during the first 1000 days is an important period to prevent undernutrition in infants and children WHO (2013). It is evidenced that suboptimal child-feeding practices are among the leading causes of child undernutrition (Tafese *et al.*, 2020). Optimal complementary feeding has been shown to reduce mortality in infants under five by 6%, according to a prior study (Black *et al.*, 2013; Oot *et al.*, 2015). Generally, undernutrition is more prevalent in nations with lower feeding rates (Disha *et al.*, 2012). Worldwide estimates of the proportion of MAD were (21%) (Lutter *et al.*, 2011). Previous works address that just 8% of children (6–23 months) in two rural districts of Pakistan had met the MAD (Khan *et al.*, 2017). Other studies from India, Nepal, and Nigeria also reported 33%, 34%, and 31.5% proportion of MDD respectively (Dhami *et al.*, 2019; Joshi *et al.*, 2012); showing the

proportion of MAD slightly varied across regions ranging from 4% in western regions of India to 34% in Nepal. Likewise, the Ethiopian Demographic and Health Survey (EDHS) 2016 reported a 7% proportion of MAD (Shagaro *et al.*, 2021).

Regional variations in the proportion of MAD were also reported in Ethiopia. Accordingly, study reports from Oromia, Afar, and southern Ethiopia reported 29%, 19.4%, and 12.3% respectively (Fentaw *et al.*, 2020; Kassa *et al.*, 2016; Wonda *et al.*, 2017). Studies from various databases published until July 2018 revealed that in Ethiopia the pooled proportion estimate of MDD, MMF, and MAD was 18.0%, (Udoh & Amodu, 2016) 56.0%, and 10.0% respectively (Abdurahman *et al.*, 2019). Studies on child feeding practices have shown that a number of factors are associated with appropriate complementary feeding practices. These include the mother's and father's educational and occupational status (Birie *et al.*, 2021; Debela *et al.*, 2021; Fentaw *et al.*, 2020; Manzione *et al.*, 2019; Mya *et al.*, 2019; Nankinga *et al.*, 2019), the family's socioeconomic status (Solomon *et al.*, 2017; Zebadia *et al.*, 2021), maternal decision-making at the household level (Beyene *et al.*, 2015; Victor *et al.*, 2014), institutional delivery (Birie *et al.*, 2021; Issaka *et al.*, 2015; Mulat *et al.*, 2019; Temesgen *et al.*, 2018), the child's age (Guirindola *et al.*, 2018; Issaka *et al.*, 2015; Mokori *et al.*, 2017; Aguayo *et al.*, 2017), receiving IYCF counseling (Liben *et al.*,


2017; Molla *et al.*, 2021; Pokhrel *et al.*, 2016), and a history of illness two weeks prior to the survey (Molla *et al.*, 2021).

Therefore, in addition to providing a better understanding of the status of child-feeding practices, the current study may produce up-to-date information on these practices and related factors that will help in the development of suitable interventions for improved infant and young child feeding (IYCF) in the study area.

## METHODOLOGY

### Study area

The study was carried out in Jigjiga City which is the capital city of the Somali regional state, located in south-eastern Ethiopia, 632 km away from Addis Ababa (Figure 1). Jigjiga experiences variation in temperature throughout the year, with highs and lows between 25.6°C and 28.2°C and 7.1°C and 15.5°C, respectively, due to its diverse weather patterns. According to the Central Statistics Agency of Ethiopia, Jigjiga has an estimated total population of 184,660 (the annual growth rate is 2.6%) of whom 46.7% are females and 53.3% are males. Among the total population, 47% are under the age of 15 years whereas 3.2 are under the age of 1 year. The reproductive age group is 22.8% and the total fertility rate (TFR) is 6.6. Life expectancy at birth is estimated to be 55.4 years for females and 58.7 years for males (Macro, 2006).



**Figure 1:** Map of the study area in Jigjiga City Somali Region, 2021, Taken from Central Statistics Agency, 2008 with slight modification.

### Study period and design

A community-based cross-sectional study design was conducted from June 15 to July 20, 2021.

### Study Population

The study populations were children aged 6 to 23 months and their mothers /caregivers (mother-child pair) who were inhabitants of the Jigjiga City administration. Respondents who were mentally or

physically disabled such as mothers who are unable to speak and/or listen, mothers with serious illness at the time of data were excluded from the study.


**Sampling Size Determination**

The single population proportion formula was used to calculate the sample size with the assumption of a 22.2% proportion of MAD in Ethiopia (Roba, O’Connor, Belachew, & O’Brien, 2016), 95% confidence level, and 5% margin of error. Adding a 10% non-response rate, the final sample size of 292 child-mother pairs was determined. Because this is a sample proportion we expect the results to be. So we have used a result from a previous survey done in most similar socioeconomic areas compared with our study area.

we also considered that the sample size calculation uses the Normal approximation to the Binomial distribution. And if the sample proportion is close to 1 as reported by EDHS (2011), (2016), and (2019) we believe the approximation cannot be valid and the sample size we get will be too small. Hence we consider an alternative proportion to get adequate power for the present study.

**Sampling Procedure**

Jigjiga City Administration has 20 kebeles, 6 kebeles were selected randomly and the total number of mothers with children aged 6-23 months was obtained from the kebele office of the selected 6 kebeles. And lastly, the study participants were selected using systematic random sampling.



**Figure 2: Schematic presentation of sampling technique of a study on minimum acceptable and associated factors in Jigjiga City Administration, Ethiopia, 2021**

**Variables (dependent and independent)**

The dependent variable was minimum acceptable diet (MAD) with the possibility of adequate or inadequate intakes whereas the independent Variables were Socio-demographic and socio-economic factors like age, marital status, family size, level of education, occupation, and income level, maternal knowledge on recommended child feeding, maternal and child-related factors (maternal and childhood illness, maternal workload), cultural norms and beliefs (presence of prohibited foods for infants, familial and cultural acceptance), and obstetric factors (Antenatal care (ANC), postnatal care (PNC), birth order, parity, place of delivery).

**Operational definitions**

**Minimum Acceptable Diet (MAD):** Proportion of children 6–23 months of age who receive a minimum acceptable diet (including breast milk). This composite indicator was calculated from children 6–23 months of age who had at least the minimum dietary diversity and the minimum meal frequency during the previous day WHO (2010).

**Minimum Dietary Diversity (MDD):** at least 5 food groups out of 8 food groups within 24 h and the cut of was <5 food is inadequate and >5 to 8 food groups consumed are adequate (Breast milk, grain, legumes, dairy products, egg, meat, fruits, and vegetables) recommended by world health organization WHO (2017).

**Minimum Meal Frequency (MMF):** cut of point of a child who receives solid, semi-solid, or soft foods (but also includes milk for non-breastfed children) the minimum number of times or more over the previous day for more the minimum number of times is 2 times for breastfed infants 6-8 months 3 times for breastfed children 9-23 months 4 times for non-breastfed children 6-23 month WHO (2010).

**Feeding Practices (FP):** these were the actions carried out by a caregiver relating to breastfeeding and complementary feeding, solid and semi-solid foods, given to a child WHO (2010).

#### **Data collection instrument**

A pre-tested structured questionnaire was used which was developed from consistent pieces of literature. The English version questionnaire was translated into the Somali language and translated back to English by an expert fluent in English and the local languages to check the consistency. The MAD was collected as a simple count of 8 food groups (breast milk, grains, roots, and tubers; legumes and nuts; dairy products; flesh foods (meats/fish/poultry); eggs; vitamin A-rich fruits and vegetables, and other fruits and vegetables) consumed by the child in 6-23 months to calculate MDD and MMF over the past 24 hours of the survey WHO (2017).

#### **Data Processing and Analysis**

The completeness and consistency of the data were checked the data was coded and double-entered into EPI Data 3.1. The data was exported into SPSS version 25 statistical software for analysis. Bivariate logistic regression was used (p-value less than 0.25) as the candidate to see the association between each independent variable and the outcome variable. Multivariable logistic regression analysis was used to control for all possible confounders and to identify independent predictors of minimum acceptable diet. The odds ratio along with a 95% confidence interval was estimated to measure the strength of the association. The level of statistical significance was declared at a P-value less than 0.05. The model

was checked for their fitness using the Hosmer and Lemeshow goodness of Fitness test.

To assure the quality of the data, the structured and pre-tested questionnaire was used. A pre-test of the questionnaire was employed before the actual data collection time among 5% of the study sample on one kebele not included in this study. During the pre-test, the questionnaire was assessed for its clarity/understandability, reliability, sensitivity of the subject matter, and cultural acceptability in the area. It was also a good opportunity to learn different techniques related to our study and, to estimate resources and time. A two-day intensive training was given on the instruments, method of data collection, and ethical issues to the data collectors.

#### **Ethical consideration**

Ethical clearance was obtained from Jigjiga University Research and Ethical Committee. An official letter was written from the school of Graduate Studies to the Jigjiga city administration to get permission and a support letter to each respected kebeles. Informed voluntary consent was sought from each participant. For illiterate women, a right thumbprint was taken as a signature. The purpose of the study was explained to the study participants. Confidentiality of responses was also ensured throughout the research process.

## **RESULTS**

### **Socio-demographic and health-related characteristics of mothers**

A total of 292 mothers were participated in the study with a response rate of 100%. The majority of participant mothers (48%) were in the age range of 26-35 years. Most of them (95.2%) were biological mothers while the rest were caregivers and family members. Nearly half of the mothers (49%) were unable to read or write, 87% were married, 95% were Muslims, and 71.2% were housewives (Table 1).

**Table 1** Socio-demographic and health care related characteristics of respondent mothers among children 6- to 23-mo-old from Jigjiga City Administration, Ethiopia 2021(n = 292)

Participant characteristic (s)	Frequency (n)	Percentage (%)
<b>Relationship to child</b>		
Biological mother	278	95.2
Caregiver (family member)	14	4.8
<b>Age of the mother(year)</b>		
15-25	84	28.8
26-35	141	48.3
36-45	62	21.2
≥45	5	1.7
<b>Marital Status</b>		
Married	254	87.0
Widowed	13	4.5
Divorced	23	7.9
Single	2	.7
<b>Family size</b>		
1-4	133	45.6
4-6	90	30.8
≥6	69	23.6
<b>Religion</b>		
Muslim	277	94.9
Christian	15	5.1
<b>Educational Status of the mothers</b>		
No formal Education	143	49.0
Primary Education (1-8 <sup>th</sup> grade)	66	22.6
High school and above	83	28.4
<b>Educational Status of the Father</b>		
No formal Education	81	27.7
Primary School (1-8 <sup>th</sup> grade)	39	13.4
High school and above	172	58.9
<b>Occupation of the mother</b>		
Housewife	208	71.2
Government employed	41	14.0
Merchant	37	12.7
Others	6	2.1
<b>Occupation of the father</b>		
Unemployed	47	16.1
Government employed	101	34.6
Merchant	94	32.2
Farmer	22	7.5
Other employment	28	9.6
<b>Total number of births</b>		
1-5	182	62.3
5-10	104	35.6
≥10	6	2.1
Mother delivered at health institution	174	59.6
<b>Sources of information for IYCF</b>		
Health care workers	169	57.9
Mass media	100	34.2
Neighbors/friends	19	6.5
Other sources	4	1.4
Mother had an ANC visit	251	86.0
Mother had a PNC visit	106	36.3
Mother ever used contraceptive methods	269	92.12

n = 292, Child, child 6–23 months old enrolled for the study; ANC: antenatal care; PNC: Post-natal care.

More than half (62.3%) of mothers gave birth to 1-5 children and 59.6% delivered at the health facility. In this study, nearly half of the mothers (57%) got IYCF information from health professionals and 50% got an extra meal during pregnancy. The majority of them (86%) had ANC and one-third (36%) of them got postnatal Care (Table 1).

### Child feeding and health-related characteristics

About 42% of the children were in the age category of 6-8 months, 56.6% were female and all children were breastfeeding. From the reported illness nearly one-third (31.5%), had diarrhea and 37% fever in the last two weeks of the survey. One hundred thirty-two children (45.2%) were fed with a cup and spoon (Table 2) and 22% of participant children met the MAD (Figure 3).

**Table 2:** Child feeding and health care characteristics among children 6- to 23-mo-old from Jigjiga City Administration, Ethiopia 2021 (n=292)

Participant characteristic (s)	N	Percentage (%)
<b>The child's age in months</b>		
6-8	123	42.1
9-11	78	26.7
12-17	58	19.9
18-23	33	11.3
<b>Gender of the child</b>		
Female	174	59.6
Male	118	40.4
Currently Breast Feeding	292	100
<b>Breastfeed initiation</b>		
Within 1 hour	108	55.4
>1 hour	87	44.6
<b>Complementary feeding initiation time</b>		
Before 6 months of age	81	27.7
At 6 months	137	46.9
After 6 months	69	23.6
Don't know	5	1.7
Child had separate plate for feeding	103	35.3
Child met minimum acceptable diet	64	22
Mother received IYCF counseling	171	58.6
Updated vaccination Status	248	84.9
Child had diarrhea in the last 2 wks	92	31.5
Child had fever in the last 2 wks	108	37.0

### Minimum acceptable diet practice Proportion

From total (292) children aged 6–23 months, about 64(22%) of

children meet the recommended minimum acceptable diet.




Figure 3: Proportion of minimum acceptable diet of children aged 6-23 months in Jigjiga city administration, Ethiopia 2021(n=292)




Figure 4: Food groups consumed among children aged 6-23 months in Jigjiga city administration, Ethiopia

**Factors associated with the minimum acceptable diet**

The binary logistic regression analysis revealed that an employed mother, a father with a higher level of education, decreased family size, less than five total number of births, institutional delivery, increased child's age, being male gender, and a mother who received IYCF counseling were found to be positively associated with MAD and significant candidates for the multivariable model (P<0.25). However, in the multivariate logistic regression analysis, only three variables were found to be significant with minimum

acceptable diet: the place of delivery, the child's age (months), and receiving IYCF counseling. Children delivered at health facilities showed a nearly two-fold higher likelihood of meeting MAD than their counterparts [AOR=1.99, 95% CI (1.07-3.70)]. Children whose age is between 12–17, and 18-23 months were 2.6 and 3.2 times more likely to meet MAD than 6–8-month children [AOR: 2.61(1.15-5.90)] and [AOR: 3.2(1.25-8.17)], respectively. The odds of MAD among mothers with children aged 6-23 months who received IYCF counseling were 2.54 times more likely than those who did not [AOR=2.54, 95% CI (1.21-5.35)] (Table 3).

**Table 3** Factors associated with minimum acceptable diet among children 6 to 23 months old from Jigjiga City Administration, Ethiopia 2021 (n=292)

Variable (s)	MAD		COR (CI95%)	AOR (CI95%)
	Meet MAD, N	Not meet MAD, N		
<b>Occupation of mother</b>				
Housewife	158(76.0)	50(24.0)	1.58(0.821-3.049)*	0.78(0.371-1.621)
Employed	70(83.3)	14(16.7)	1	1
<b>Educational status of Father</b>				
Formal	68(84.0)	13(16.0)	1.67(0.852-3.264)*	1.07(0.490-2.324)
Informal	160(75.8)	51(24.2)	1	1
<b>Family Size, n</b>				
<=5	96(72.2)	37(27.8)	0.53(0.303-0.930)*	0.55(0.284-1.046)
>5	132(83.0)	27(17.0)	1	1
<b>Total births, n</b>				
1-5	138(75.8)	44(24.2)	1	1
>5	90(81.8)	20(18.2)	0.77(0.386-1.259)*	0.74(0.379-1.433)
<b>Institutional delivery</b>				
Yes	86(72.9)	32(27.1)	1.65 (0.945-2.886)**	1.99(1.069-3.704)**
No	142(81.6)	32(18.4)	1	1
<b>Age of the child, mo</b>				
9–11	63(80.8)	15(19.2)	1.23 (0.585-2.568)*	1.39(0.625-3.081)
12–17	41(70.7)	17(29.3)	2.14 (1.018-4.480)**	2.61(1.152-5.92)**
18–23	21(63.6)	12(36.4)	2.94 (1.251-6.925)**	3.2(1.246-8.168)**
6–8	103(83.7)	20(16.3)	1	1
<b>Gender of the child</b>				
Male	114(96.6%)	4(3.4%)	0.07 (0.77-0.19)*	1.99(1.069-3.704)
Female	114(65.5%)	60(34.5%)	1	1
<b>Mother ever had IYCF counseling</b>				
Yes	121(70.3)	51(29.7)	3.5(1.789-6.727)**	2.54(1.210-5.348)**
No	107(89.2)	13(10.8)	1	1

Notice \* p-value < 0.25(COR- crude odds ratio), \*\*p-value < 0.05, (AOR- adjusted odds ratio).

## DISCUSSION

This study has assessed the proportion of MAD and associated factors among children aged 6–23 months in Jigjiga City, Somali region of Ethiopia. The proportion of children who met the MAD in the present study was 22% (n = 64). These findings are consistent with studies conducted in Ethiopia and elsewhere which was 21% from the estimates of surveys of 147 countries (Lutter *et al.*, 2011). The percentage of children met MAD in Nepal were 26.5% (Joshi *et al.*, 2012) and 19.4% in Afar Region of Ethiopia (Fentaw *et al.*, 2020). This result is lower compared with the result reported from studies conducted in Congo 33% (Kambale *et al.*, 2021), Addis Ababa, Ethiopia 74.6% (Abebe *et al.*, 2021), Mareka District, Southern Ethiopia 35.5% (Feleke & Mulaw, 2020), and Slums of Oromia Zones 29% (Wondu Garoma & Yang, 2017). But this result is higher compared to the report of the Ethiopian demographic and health survey 11% (Shagaro *et al.*, 2021), Pakistan 8% (Khan *et al.*, 2017), India 7.3% (Dhami *et al.*, 2019), Nigeria 7.3% (Udoh & Amodu, 2016). This difference may be due to differences in study settings, and socio-demographic and economic characteristics of

the study participants and might also be due to sampling size, study area, and period.

The present study identified factors associated with MAD that are consistent with similar studies done across Ethiopia and elsewhere. Place of delivery is one of the identified factors significantly associated with MAD showing that children born at the health facility are positively associated with meeting the MAD than their counterparts. A similar finding was reported by prior studies conducted in Ghana (Issaka *et al.*, 2015), and North West Ethiopia (Birie *et al.*, 2021; Mulat *et al.*, 2019; Temesgen *et al.*, 2018). This may be related to the fact that those mothers delivered in the health facility may get nutritional counseling by the health professionals and this can persuade them to a favorable attitude toward appropriate complementary feeding practice.

The other significant factor associated with the child feeding practice was the child's age. Those children aged one year and above showed more than double-fold likelihood to meet MAD as compared to those under one year of age. This finding is similar to the studies conducted in central Ethiopia (Molla *et al.*, 2021), the Philippines (Guirindola *et al.*, 2018), Uganda (Mokori *et al.*, 2017), Ghana (Issaka *et al.*, 2015), and Pakistan (Khan *et al.*, 2017).



Mothers (caregivers) in Ethiopia often complain about the lack of sufficient options for local complementary foods for a child to feed during the early age (under 1 year) of the complementary period. Commercially produced complementary foods are beyond the reach of most households (Ayele *et al.*, 2022). The odds of MAD among children whose mothers received IYCF counseling were more than twofold more likely to meet MAD than those who did not. This result is in line with studies conducted in Ethiopia (Liben *et al.*, 2017; Molla *et al.*, 2021) and Nepal (Karn *et al.*, 2019; Pokhrel *et al.*, 2016). The positive association of maternal exposure to IYCF counseling could be explained in terms of the role of child-feeding counseling for the mother in increasing her knowledge and changing her attitude favorably towards appropriate child-feeding practice and this may play an important role in meeting the standard of MAD for her child.

## CONCLUSION

The study revealed that the children delivered in healthcare facilities are more likely to meet the recommended MAD compared to those delivered at home. This could be due to the availability of trained healthcare professionals and access to proper nutrition support using healthcare facilities. Children who received counseling on infant and young child feeding (IYCF) were more likely to meet the recommended MAD. This highlights the importance of educating caregivers on appropriate feeding practices, which can positively impact children's nutrition. The specific age group within the 6-23 month range also influences the likelihood of meeting the recommended MAD. Further analysis is needed to understand the specific factors that affect different age groups within this range. Efforts should be made to improve access to healthcare facilities for deliveries, especially in areas with limited healthcare infrastructure. This can ensure that the children would have access to trained professionals and proper nutrition support from birth. Investing in IYCF counseling programs is crucial to improve caregivers' knowledge and practices related to infant and young child feeding. These programs should be implemented at various levels, including antenatal care, postnatal care, and community-based settings. As different age groups (within the 6-23-month range) have varying factors influencing their dietary practices, interventions should be tailored accordingly. Further research should be conducted to identify specific barriers and facilitators for each age group, and targeted interventions should be developed to address these factors. Overall, addressing these factors can contribute to improving the proportion of children meeting the recommended MAD and ultimately enhance the nutritional status and well-being of children aged 6-23 months.

## Strength

The fact that this study was carried out at the community level raises the likelihood that its findings will be applicable to the complete population from which the sample was taken. To account for all potential confounders, a multivariate logistic regression analysis was utilized in this investigation.

## Limitations

The study only used a 24-hour recall method for food intake, so it's possible that the results might not correctly reflect participants' past feeding dietary habits. Additionally, there could be recall bias, and since this was a self-reported study, the minimum amount of dietary variety practice might not have been accurately recorded (social desirability bias).

## ACKNOWLEDGMENTS

The authors are delighted to acknowledge Jigjiga University for the ethical clearance and any other support to this research. We also give thanks to all individuals who participated in this study particularly mothers who gave their time in this study and the data collectors who diligently worked to complete this study.

## CONFLICTS OF INTEREST

The authors declare that they have no competing interests.

## FUNDING

The authors have not declared a specific grant for this research from any funding agency in the public, commercial, or not-for-profit sectors.

## CONTRIBUTIONS

All authors made a significant contribution to the present study. AAI conceptualized the study, developed the study design, and was involved in the analysis and write-up. ZT and AMA were involved in the analysis, writing up, and interpretation of the data. AA and SMH wrote the first draft. SMH and ZA reviewed and edited the draft manuscript. All authors have reviewed the final manuscript and approved it before submission.

## DATA AVAILABILITY

The data for this study are available from the first author upon reasonable request.

## CONSENT FOR PUBLICATION

Not applicable.

## REFERENCES

- Abdurahman, A. A., Chaka, E. E., Bule, M. H., & Niaz, K. (2019). Magnitude and determinants of complementary feeding practices in Ethiopia: A systematic review and meta-analysis. *Heliyon*, 5(7).
- Abebe, H., Gashu, M., Kebede, A., Abata, H., Yeshaneh, A., Workye, H., & Adane, D. (2021). Minimum acceptable diet and associated factors among children aged 6–23 months in Ethiopia. *Italian Journal of Pediatrics*, 47(1), 1-10.
- Ayele, D. A., Teferra, T. F., Frank, J., & Gebremedhin, S. (2022). Optimization of nutritional and functional qualities of local complementary foods of southern Ethiopia using a customized mixture design. *Food Science & Nutrition*, 10(1), 239-252.

- Beyene, M., Worku, A. G., & Wassie, M. M. (2015). Dietary diversity, meal frequency and associated factors among infant and young children in Northwest Ethiopia: a cross-sectional study. *BMC Public Health*, 15(1), 1-9.
- Birie, B., Kassa, A., Kebede, E., & Terefe, B. (2021). Minimum acceptable diet practice and its associated factors among children aged 6–23 months in rural communities of Goncha district, north West Ethiopia. *BMC nutrition*, 7(1), 1-9.
- Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., De Onis, M., . . . Martorell, R. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries. *The lancet*, 382(9890), 427-451.
- Debela, B. L., Gehrke, E., & Qaim, M. (2021). Links between maternal employment and child nutrition in rural Tanzania. *American Journal of Agricultural Economics*, 103(3), 812-830.
- Dhami, M. V., Ogbo, F. A., Osuagwu, U. L., & Agho, K. E. (2019). Prevalence and factors associated with complementary feeding practices among children aged 6–23 months in India: a regional analysis. *BMC Public Health*, 19(1), 1-16.
- Disha, A., Rawat, R., Subandoro, A., & Menon, P. (2012). Infant and young child feeding (IYCF) practices in Ethiopia and Zambia and their association with child nutrition: analysis of demographic and health survey data. *African Journal of Food, Agriculture, Nutrition and Development*, 12(2), 5895-5914.
- Feleke, F. W., & Mulaw, G. F. (2020). Minimum acceptable diet and its predictors among children aged 6-23 months in Mareka District, southern Ethiopia: community based cross-sectional study. *International Journal*, 9(4), 203.
- Fentaw Mulaw, G., Wassie Feleke, F., & Adane Masresha, S. (2020). Maternal characteristics are associated with child dietary diversity score, in Golina District, Northeast Ethiopia: a community-based cross-sectional study. *Journal of Nutrition and Metabolism*, 2020.
- Guirindola, M. O., Maniego, M. L. V., Silvestre, C. J., & Acuin, C. C. S. (2018). Determinants of meeting the minimum acceptable diet among Filipino children aged 6-23 months. *Philipp J Sci*, 147(1), 75-89.
- Issaka, A. I., Agho, K. E., Burns, P., Page, A., & Dibley, M. J. (2015). Determinants of inadequate complementary feeding practices among children aged 6–23 months in Ghana. *Public Health Nutrition*, 18(4), 669-678.
- Joshi, N., Agho, K. E., Dibley, M. J., Senarath, U., & Tiwari, K. (2012). Determinants of inappropriate complementary feeding practices in young children in Nepal: secondary data analysis of Demographic and Health Survey 2006. *Maternal & Child Nutrition*, 8, 45-59.
- Kambale, R. M., Ngaboyeka, G. A., Kasengi, J. B., Niyitegeka, S., Cinkenye, B. R., Baruti, A., . . . Van der Linden, D. (2021). Minimum acceptable diet among children aged 6–23 months in South Kivu, Democratic Republic of Congo: a community-based cross-sectional study. *BMC Pediatrics*, 21(1), 1-9.
- Karn, S., Adhikari, D. P., Paudyal, N., Aryal, B., Adhikari, R. K., & Steffen, M. M. (2019). Child undernutrition and feeding practices in Nepal: trends, inequities, and determinants. *DHS Further Analysis Report(122)*.
- Kassa, T., Meshesha, B., Haji, Y., & Ebrahim, J. (2016). Appropriate complementary feeding practices and associated factors among mothers of children age 6–23 months in Southern Ethiopia, 2015. *BMC Pediatrics*, 16(1), 1-10.
- Khan, G. N., Ariff, S., Khan, U., Habib, A., Umer, M., Suhag, Z., . . . Turab, A. (2017). Determinants of infant and young child feeding practices by mothers in two rural districts of Sindh, Pakistan: a cross-sectional survey. *International Breastfeeding Journal*, 12, 1-8.
- Liben, M. L., Abuhay, T., & Haile, Y. (2017). Factors associated with dietary diversity among children of agro pastoral households in afar regional state, northeastern Ethiopia. *Acad. J. Ped. Neonatol*, 5(5).
- Lutter, C. K., Daelmans, B. M., de Onis, M., Kothari, M. T., Ruel, M. T., Arimond, M., . . . Borghi, E. (2011). Undernutrition, poor feeding practices, and low coverage of key nutrition interventions. *Pediatrics*, 128(6), e1418-e1427.
- Macro, O. (2006). Central Statistical Agency Addis Ababa, Ethiopia. Central Statistical Agency Addis Ababa, Ethiopia.
- Manziona, L. C., Kriser, H., Gamboa, E. G., Hanson, C. M., Mulokozi, G., Mwaipape, O., . . . Hall, P. C. (2019). Maternal employment status and minimum meal frequency in children 6-23 months in Tanzania. *International Journal of Environmental Research and Public Health*, 16(7), 1137.
- Mokori, A., Schonfeldt, H., & Hendriks, S. L. (2017). Child factors associated with complementary feeding practices in Uganda. *South African Journal of Clinical Nutrition*, 30(1), 7-14.
- Molla, A., Egata, G., Getacher, L., Kebede, B., Sayih, A., Arega, M., & Bante, A. (2021). Minimum acceptable diet and associated factors among infants and young children aged 6–23 months in Amhara region, Central Ethiopia: community-based cross-sectional study. *BMJ open*, 11(5), e044284.
- Moore, T., Arefadib, N., Deery, A., West, S., & Keyes, M. (2017). The first thousand days: An evidence paper-summary.
- Mulat, E., Alem, G., Woyraw, W., & Temesgen, H. (2019). Uptake of minimum acceptable diet among children aged 6–23 months in orthodox religion followers during fasting

- season in rural area, DEMBECHA, north West Ethiopia. *BMC Nutrition*, 5, 1-10.
- Mya, K. S., Kyaw, A. T., & Tun, T. (2019). Feeding practices and nutritional status of children age 6-23 months in Myanmar: A secondary analysis of the 2015-16 Demographic and Health Survey. *PLoS one*, 14(1), e0209044.
- Na, M., Aguayo, V. M., Arimond, M., & Stewart, C. P. (2017). Risk factors of poor complementary feeding practices in Pakistani children aged 6–23 months: A multilevel analysis of the Demographic and Health Survey 2012–2013. *Maternal & Child Nutrition*, 13, e12463.
- Nankinga, O., Kwagala, B., & Walakira, E. J. (2019). Maternal employment and child nutritional status in Uganda. *PLoS one*, 14(12), e0226720.
- Ng, C. S., Dibley, M. J., & Agho, K. E. (2012). Complementary feeding indicators and determinants of poor feeding practices in Indonesia: a secondary analysis of 2007 Demographic and Health Survey data. *Public Health Nutrition*, 15(5), 827-839.
- Oot, L., Sommerfelt, E., Sethuraman, K., & Rossimize, J. (2015). Estimating the effect of suboptimal breastfeeding practices on child mortality: A model in profiles for country-level advocacy. Technical brief. Food and Nutrition Technical Assistance III Project, 1-9.
- WHO (2010). Indicators for assessing infant and young child feeding practices: part 2: measurement: World Health Organization.
- WHO (2013). Essential nutrition actions: improving maternal, newborn, infant and young child health and nutrition.
- WHO (2017). Global Nutrition Monitoring Framework: operational guidance for tracking progress in meeting targets for 2025.
- WHO (2018). Nurturing care for early childhood development: a framework for helping children survive and thrive to transform health and human potential.
- Pokhrel, K., Nanishi, K., Poudel, K. C., Pokhrel, K. G., Tiwari, K., & Jimba, M. (2016). Undernutrition among infants and children in Nepal: maternal health services and their roles to prevent it. *Maternal and Child Health Journal*, 20, 2037-2049.
- Prado, E. L., & Dewey, K. G. (2014). Nutrition and brain development in early life. *Nutrition Reviews*, 72(4), 267-284.
- Roba, K. T., O'Connor, T. P., Belachew, T., & O'Brien, N. M. (2016). Infant and young child feeding (IYCF) practices among mothers of children aged 6–23 months in two agro-ecological zones of rural Ethiopia. *Int J Nutr Food Sci*, 5(3), 185-194.
- Shagaro, S. S., Mulugeta, B. e. T., & Kale, T. D. (2021). Complementary feeding practices and associated factors among mothers of children aged 6-23 months in Ethiopia: Secondary data analysis of Ethiopian mini demographic and health survey 2019. *Archives of Public Health*, 79(1), 205. doi:10.1186/s13690-021-00725-x
- Solomon, D., Aderaw, Z., & Tegegne, T. K. (2017). Minimum dietary diversity and associated factors among children aged 6–23 months in Addis Ababa, Ethiopia. *International Journal for Equity in Health*, 16(1), 1-9.
- Tafese, Z., Reta Alemayehu, F., Anato, A., Berhan, Y., & Stoecker, B. J. (2020). Child feeding practice and primary health care as major correlates of stunting and underweight among 6-to 23-month-old infants and young children in food-insecure households in Ethiopia. *Current Developments in Nutrition*, 4(9), nzaa137.
- Temesgen, H., Yeneabat, T., & Teshome, M. (2018). Dietary diversity and associated factors among children aged 6–23 months in Sinan Woreda, Northwest Ethiopia: a cross-sectional study. *BMC Nutrition*, 4(1), 1-8.
- Udoh, E. E., & Amodu, O. K. (2016). Complementary feeding practices among mothers and nutritional status of infants in Akpabuyo Area, Cross River State Nigeria. *SpringerPlus*, 5, 1-19.
- Victor, R., Baines, S. K., Agho, K. E., & Dibley, M. J. (2014). Factors associated with inappropriate complementary feeding practices among children aged 6–23 months in Tanzania. *Maternal & Child Nutrition*, 10(4), 545-561.
- Wondu Garoma, B., & Yang, N. (2017). Determinants of suboptimal complementary feeding practices among children aged 6-23 months in selected urban slums of Oromia zones (Ethiopia). *J Nutr Food Sci*, 7(593), 2-13.
- Zebadia, E., Mahmudiono, T., Atmaka, D. R., Dewi, M., Helmyati, S., & Yuniar, C. T. (2021). Factors Associated with Minimum Acceptable Diet in 6–11-Month-Old Indonesian Children Using the 2017 IDHS. Open access Macedonian *Journal of Medical Sciences*, 9(E), 1403-1412.