

Nutrient Release Pattern from Leptic Cambisols as Influenced by Vermicompost and Inorganic Fertilizer Applications

Molla Hadis^{1*}, Gashaw Meteke² and Wassie Haile²

¹Tigray Agricultural Research Institute, Mekelle Soil Research Center, Mekelle, Ethiopia

²School of Plant and Horticultural Science, College of Agriculture, Hawassa University,

P. O. Box 05, Hawassa Ethiopia

Abstract	Article Information
<p>An experiment was conducted to determine the effects of vermicompost, inorganic fertilizers and their combinations on release of soil nutrients, at different growth stages of wheat. A factorial combination of four levels of inorganic fertilizers (33.33 %, 66.66%, and 100 % of the recommended NPK fertilizers) and vermicompost (0, 2, 4 and 6 t ha⁻¹) 0, were laid out in Complete Randomized Design (CRD) with three replications. Soil was collected before planting and after planting (at tillering, flowering and maturity stages of wheat) from each pot in order to determine dynamics of selected nutrients [Nitrogen (N), Phosphorus (P) and Potassium (K)]. The interaction between vermicompost and chemical fertilizers were not significant for NPK contents of the soil at all growth stages except phosphorus at heading stage. In all cases, highly significant increases in total N, available P and K in the soil were observed due to the increasing rates of main effect vermicompost or inorganic fertilizers during all growing periods. The highest available as well as total contents of NPK in the soil were found at tillering stage. This initial increment at tillering stage for both factors showed a declining trend later at heading and maturity stages. However, the observed decline was in exception for vermicompost applied at 6 t ha⁻¹ which maintained highest level of available P and K and 4 t ha⁻¹ which continued mineralization of K up to heading stage. In general, application of 6 tone vermicompost per hectare was found proportional with the full dose of the recommended fertilizers in supplying NPK for wheat crop. Therefore, building up the total as well as available NPK to higher levels up to heading stage can bring maximum nutrient uptake and yields of wheat.</p>	<p>Article History: Received : 10-01-2016 Revised : 16-03-2016 Accepted : 23-03-2016</p> <hr/> <p>Keywords: Nutrient availability, growth stages, mineralization, nutrient decline</p> <hr/> <p>*Corresponding Author: Mola Hadis¹ E-mail: moheaddis@gmail.com</p>

Copyright©2016 STAR Journal, Wollega University. All Rights Reserved.

INTRODUCTION

In Ethiopia, the major constraint to agricultural growth and food self-sufficiency for a long period is the decline in soil fertility. Soil fertility

is a manageable soil property and its management is of utmost importance for optimizing crop nutrition on both a short-term

and a long-term basis to achieve sustainable crop production (Roy *et al.*, 2006). In response to this problem, major efforts have been made to encourage the wider use of inorganic fertilizers. Regardless of a five times increase in fertilizer application in Ethiopia, national cereal yields have only increased 10 percent since the 1980s (Gete *et al.*, 2010), while relative benefits of chemical fertilizer application have decreased over time.

Even though chemical fertilizers substantially increased the available plant nutrients during the first weeks, a losses arising from different factors limits the continual supply of nutrients at critical periods where plants inquire high demand for these nutrients. Reports from Hammermeister *et al.* (2006) show that chemically fertilized plots declined in NH_4^+ as sampling time continues. In other hand, available P was high at tillering which declined later at flowering and maturity stages of rice (Lungmuana *et al.*, 2013) compared to organic residues and manures. Loss of ammonium from Urea due to leaching and volatilization during two to three weeks after application (Jones *et al.*, 2007), fixation of phosphorus with Al/Fe oxides (Lungmuana *et al.*, 2013) and potassium with interlayer of clay mineralogy (Najafi-ghiri, 2014) were reported after the addition of inorganic fertilizers. Such losses could exhaust the pool and available stock of certain nutrients and brought a question whether they could support plant life until the maturity stage.

However, organic farming systems with the aid of various nutrients of biological origin such as compost and vermicompost were thought to be the answers for the 'food safety and farm security' in future (Sinha *et al.*, 2009). On the other hand, initially low levels and low release of available nutrients from organic amendments at seedling and vegetative stages can restrict uptake of adequate nutrients that might result poor root and shoot growth of plants. Additionally, the ill effects of farmlands brought about by the application of chemical agricultural inputs for long periods in favor of boosting plant

yields (Arancon *et al.*, 2006; Sheoran *et al.*, 2015; and Sinha *et al.*, 2009) were at a cost of inherent soil fertility and microbe inhabitants; which are supposed to maintain the balance of rhizosphere by natural law.

Furthermore, NPK deficiencies are widespread and external applications are necessary to augment soil supplies for harvesting optimal crop yields while minimizing the depletion of soil nutrient reserves (Roy *et al.*, 2006). In Ethiopia, it was reported that some crops have been suffering from deficiencies of nutrients other than nitrogen and phosphorus (ATA, 2014). This cannot be maintained in agricultural soils of the country where only NP fertilizers are applied unless we do not supply adequately with organic amendments like vermicasts.

Thus, evaluation of such new technologies when applied solely and/or integrated with the chemical fertilizers might give farmers another alternative to overcome such problems. At the same time building sustainable and climate smart agriculture with vermicasts can bring a new outlook to organic farming. Therefore, this research was conducted to investigate the effect of sole and combined applications of organic and inorganic fertilizers on the release of NPK at different growth stages of wheat.

Materials and Methods

Area Description

Soil for pot experiment was taken from farmlands of Mekan area, Enda-Mehoni district, Southern Tigray, Ethiopia. Soils having the same cropping history and soil types were sampled for potting media. According to the pedological map developed for the district, the soil type of the sampling area is Leptic Cambisols (Mitiku *et al.*, 2007). The sampling area is located at 39° 29'18" up to 39° 33'35" longitude and 12° 43'28" up to 12° 46'12" latitude. The pot experiment was carried out at MIT Tissue Culture Micro-Propagation Laboratory's Greenhouse, Mekelle, Ethiopia.

Treatments and Experimental Designs

Vermicompost and chemical fertilizers was used in this experiment as a source of nutrients (NPK). Vermicompost was processed by earthworm (*Eisinea fetida*) using cow manure, *Lanthana Camara* leaves and wheat straw. The soil was filled to forty-eight plastic pots having 30 and 20 cm upper and bottom diameters, respectively and 28 cm depth.

A factorial combination of four levels vermicompost and four levels inorganic fertilizers (NPK) was laid out in Complete Randomized Design (CRD) with three replications. The vermicompost levels were consisted of 0, 2, 4 and 6 t ha⁻¹ and inorganic fertilizers (NPK) were 0, 33.33 %, 66.66%, and 100 % of the recommended NPK rates. All the rates of vermicompost and NPK from inorganic fertilizers were incorporated once irrespective of the treatments and moistened at optimum. The elemental nitrogen (N), phosphorus (P) and potassium (K) was applied in the form of urea, triple superphosphate (TSP), and murate of potash (KCl) respectively. The full doses or recommended rates of NPK fertilizers are the blanket recommendation of 64 kg N ha⁻¹, 46 kg P₂O₅ ha⁻¹ which is a common practice for cereal crops throughout the country and 60 Kg K₂O ha⁻¹ respectively. The amount of Urea, TSP and KCl (g pot⁻¹) was determined by multiplying the recommended fertilizer rates (kg ha⁻¹) and 5 kg soil pot⁻¹ and divided by 2000000 kg soil ha⁻¹. Improved wheat variety; Kekaba was used as a test crop and eight seeds of wheat were planted per pot and thinned to 5 after germination. The moisture level was monitored regularly and maintained with distilled water.

Sampling and Laboratory Analysis of Soils and Vermicompost

Prior to planting, one composite soil sample from all soil sampling points and from the processed vermicompost was taken for routine analysis and the result is presented in Table 2. Soil sample was also collected after planting at tillering, flowering and ripening stages of wheat

from each pot. The collected soil sample was air-dried, milled using pistil and mortar, sieved to pass through 2mm diameter mesh sieve, stored and tagged in plastic bags. Then, particle size distribution, pH, EC, CEC, %OC, total N, available P, exchangeable K analyses was carried out for vermicompost and soil samples before planting and Total N (mg Kg⁻¹), Av.P (ppm) and Av.K (ppm) for the successive periodic soil samples to investigate the dynamics of nutrients from the above treatments. Soil texture was determined by the hydrometer method (Bouyoucos, 1962).

Soil pH and EC was determined using a pH meter from 1:2.5 soil:water ratio suspension as described in Rhoades (1982) and using EC meter from 1:5 soil water saturation extract (Jakson, 1967) respectively. The determination of soil organic carbon was based on the Walkley-Black chromic acid wet oxidation method (Walkley and Black, 1934). The Kjeldhal process (digestion, distillation and titration) as outlined by Bremner et al. (1982) was followed to determine the total nitrogen. Olsen Method (Bicarbonate extractable P) was used to extract and determine available phosphorus, using 0.5 M NaHCO₃ at adjusted pH 8.5 (Olsen et al., 1954).

Determination of CEC at pH 7 was carried out with Ammonium Acetate method as described by Chapman (1965). The amount of exchangeable cations (K) in the extract was determined by flame photometer according to Gupta (2000). Available K was determined by extracting from the sample with Morgan's solution as outlined by Sahlemedhin and Taye (2000).

RESULTS AND DISCUSSION

Selected Physicochemical Properties of the Soil and Vermicompost

The textural class of the soil used for the pot experiment was sandy clay loam. According to the rating made by Tekalign (1991), the soil reaction (pH) was moderately alkaline and the organic amendment (vermicompost) was neutral.

Table 1: Initial characteristics of the soil and vermicompost used for pot experiment

Parameters measured	Sample Source	
	Soil	Vermicompost
pH	7.48	6.78
EC (ms m ⁻¹)	0.05	2.77
CEC (cmol(+) kg ⁻¹)	30.6	-
% OC	0.98	11.37
% OM	1.68	19.60
Total N (mg Kg ⁻¹)	600	14100
% Total P	-	0.78
Av P (ppm)	9.26	-
% Total K	-	1.02
Exc.K (cmol(+) kg ⁻¹)	0.34	-
% Clay	45	-
% Silt	27	-
%Sand	28	-
Textural Class	Sandy clay loam	-
Moisture content (%)	-	38

Similarly, the total organic carbon (%) and organic matter (%) of the soil was found low, whereas it was very high for vermicompost.

The cation exchange capacity (CEC) of the soil was found to be high as outlined by Hazelton and Murphy (2007). There was medium total nitrogen (%) content on the soil and very high on the vermicompost which used as potting media according to the rating made by Berhanu (1980) and Tekalign (1991). The total and available phosphorus content of the vermicompost and the soil was rated as very high (Murphy, 1968) and medium (Cottenie, 1980) respectively. Moreover, the total and exchangeable amounts of potassium on the vermicompost

and the soil were also found to be medium (FAO, 2006), respectively.

Total nitrogen (mg kg⁻¹) content of the soil at different growth stages

The interaction between vermicompost and Inorganic fertilizers was not significant for nitrogen content of the soil at all growth stages, but there were highly significant main effects ($P \leq 0.0001$). In all cases, a highly significant increase ($P \leq 0.0001$) in total N of the soil was observed due to the increasing rates of vermicompost or inorganic fertilizers during all growing periods. These results were in agreement with the findings of Thakare and Wake (2015).




Figure 1: Total Nitrogen (mg kg⁻¹) content of the soil at different growth stages of wheat as influenced by application of different vermicompost levels




Figure 2: Total Nitrogen (mg kg^{-1}) content of the soil at different growth stages of wheat as influenced by application of different chemical fertilizer level

Out of all growth stages, the highest total N of the soil was measured at tillering stage for both main effects. At this stage, the total N was 738, 1137, 1495, 1909 mg kg^{-1} for the pots received 0, 2, 4 and 6 t VC ha^{-1} (Fig. 2) and 979, 1196, 1425, and 1679 mg kg^{-1} for the pots received 0, 33.33, 66.67 and 100 % of the recommended doses of NPK, respectively (Fig. 3). However, the total N (mg kg^{-1}) content for both main effects showed a declining trend from tillering to heading and maturity stages for all treatments (Fig. 2 & 3). In that order, it declined by 23, 19, 21 and 23 % for the rates of main effect vermicompost and about 30, 21, 17 and 21 % for the rates of main effect inorganic fertilizers (NPK) during heading stage. Accordingly, about 39, 51, 54 and 58 % of the total N measured at tillering from pots received 0, 2, 4 and 6 t VC ha^{-1} (Fig. 2), respectively were declined at maturity stage of the crop. Similarly, the declines for pots treated with 0, 33.33, 66.67 and 100 % of the recommended doses of NPK from inorganic fertilizers at this stage were 48, 51, 53 and 56 %, respectively. This trend has also been reported by Hammermeister *et al.* (2006), who reported lower soil N mineral content in the final analysis compared to the initial analysis for several treatments, suggesting loss of N with time. Similarly, Nathiya and Sanjivkumar (2015) have also found higher available nitrogen content on vegetative and flowering stage of groundnut than at post harvest stage, indicating that plants derive nutrients from soil for their growth and

development leading to the depletion of soil nutrients at the late growth stages. The decline in mineralized nitrogen with time might be related to the decrease in the labile organic matter (Tirol-Padre *et al.*, 2007).

The temporal distribution of the total nitrogen content for the main effect chemical fertilizer had the same trend with vermicompost. The total nitrogen content was increased with application levels of inorganic fertilizers at all growth stages and reached its maximum at tillering, which sharply declined at heading and maturity stages (Fig. 3). Such initial increasing trend in available N after application of high rates of N from chemical fertilizer has also been reported, which later declined significantly with sampling time from tillering to heading and maturity stages of spring wheat (Lu *et al.*, 2010).

The extent of decline in total nitrogen content from all treatment levels except the control pots was higher after heading stage of wheat for both treatment levels. At the same time, this decline was increased as the application rate of vermicompost and chemical fertilizers increased. Similar results have reported by Lu *et al.* (2010) who measured an excessive N loss due to application of high N rate. In this experiment, elevated reduction with application of high N levels could be explained due to the observed higher uptake of nitrogen. Studies from Mehta *et al.*, (1963) indicated that higher nutrient uptake at the later stages of wheat, which could serve as an

exception to the decline in the total N content of the soil solution. On the other hand, immobilization of nutrients explained by increases in the microbial biomass in the soils treated with vermicomposts could decline N contents of the soil and such increased in microbial biomass were greater in soils receiving higher rates of vermicompost applications (Arancon *et al.*, 2006).

It was observed that, the average decline in nitrogen content from the experiment was relatively higher in chemically fertilized pots than for vermicompost treatments. Moreover, the relative higher amounts of total nitrogen measured at all growth stages with application of highest doses of vermicompost (6 t ha⁻¹) compared to other vermicompost and chemical fertilizer levels might be due to the higher level of initial organic carbon content (11.4%) of the vermicompost (Table 2). Similar conclusion had been made from previous studies, which verified that changes in the organic carbon content brought about changes in the total nitrogen content of the soil (Angelova *et al.*, 2013; Tirol-Padre *et al.*, 2007). This investigation also showed a strong positive correlation between the total organic carbon content and the total nitrogen content with value of correlation coefficient as high as 0.97 (Angelova *et al.*, 2013). Angelova *et al.*, (2013) have also indicated that the considerable amount of humic acid present in vermicompost could serve as a binding site to NH₄⁺, which could prevent from the possible losses through leaching and volatilization.

Phosphorus release pattern at different growth stages

Application of vermicompost and chemical fertilizers affected the availability of phosphorus in the soil. All the experimental pots treated with vermicompost and inorganic fertilizers released higher amounts of available P over the control and the availability significantly increased at all growth stages ($p \leq 0.0001$), as of the application rate increases. Application of 2, 4 and 6 t ha⁻¹ vermicompost gave an advantage of 28%, 48% and 76% available P, respectively, over the control at tillering stages of wheat. The increase in available P due to vermicompost was in agreement with the findings of Angelova *et al.* (2013) and Tharmaraj *et al.*, (2010). Generally,

application of a considerable amount of total phosphorus (0.78% P in VC) (Table 2) with highest doses of vermicompost, could attribute to the elevated increase in available phosphorus irrespective of the treatment levels. After the initial increment in available phosphorus at tillering, a decline was observed at heading stage except for the higher doses of vermicompost (4 and 6 t ha⁻¹) (Fig. 4). The available phosphorus content for the rates of main effect vermicompost was then decreased at maturity stage, by average of 43%.

These higher doses of vermicompost continue the release of available phosphorus at tillering, maintain highest level up to heading, and then decreased at maturity by satisfying the P demand of wheat (Fig. 4). Such increase on available P at mid stages of crops (mainly at vegetative and flowering) were reported by Lungmuana *et al.* (2013), Malik *et al.* (2013) and Nathiya and Sanjivkumar (2015). The increase in available phosphorus at mid stage of wheat growth might be due to the continuous breakdown of organic matter and solubilization of phosphate minerals by soil microorganisms. According to Arancon *et al.* (2006), the increased amounts of orthophosphates in the soil from the vermicompost-treated plots could be explained by the significant correlations between the microbial biomass N and orthophosphates, indicating that release of P was due largely to the activity of soil microorganisms.

On the other hand, application of 33.33 %, 66.67 % and 100 % NPK showed 32%, 56% and 89% increments on available P contents, respectively, over the control at tillering stage. Unlike vermicompost, chemical fertilizers slightly decreased at heading stage and sharply reduced in available P at maturity stages. On average, a decline of 44 % on available P was observed for the main effect inorganic fertilizers, at maturity. However, pots treated with highest doses of chemical fertilizers (100 % NPK) have relatively higher phosphorus content (29.23 ppm) at tillering stage than vermicompost treated pots, whereas at maturity the application of highest-level vermicompost has resulted in higher available P (18.60 ppm) than did the other treatments.

The readily soluble P applied as chemical fertilizers might be exposed to P-fixation reactions in the soil, which might result in periodic decline of available P (Lungmuana *et al.*, 2013). The decline

of nutrients might also be due to the adverse effects of chemical fertilizers on beneficial soil microorganisms and soil chemistry, which would

hamper the mineralization and solubilization of P from the soil.




Figure 3: Phosphorus release pattern in the soil at different growth stages of wheat as influenced by application of different vermicompost levels




Figure 4: Phosphorus release pattern in the soil at different growth stages of wheat as influenced by application of chemical fertilizers

Potassium (K) release pattern at different growth stages

The interaction of the different levels of vermicompost and NPK fertilizer was non-significant on the release of available potassium at all growth stages. However, there were marked differences due to fertilization with either main effect vermicompost or inorganic fertilizers. The availability of K (ppm) at all growth stages notably increased ($p \leq 0.0001$) as the application rates of vermicompost and inorganic fertilizer increase

(Fig. 7 and Fig. 8). The maximum available K values for the main effect of vermicompost were 386.92, 402.42 and 288.17 ppm for pots treated with 6 t ha⁻¹, while the minimum were 289.58, 254.83 and 208.17 ppm for the control treatment at tillering, heading and maturity stages of wheat, respectively. Many studies have reported that a significantly higher values of available K obtained after the introduction of vermicompost (Angelova *et al.* 2013; Pankajam and Davi, 2009; Tharmaraj *et al.*, 2010). This increase might be

due to the application of a considerable amount of available K found in the earthworm processed organic matter (1.02%) (Table 2) and its solubilizing effect may result in release of available K from the soil.

The highest available K values for the chemical fertilizer applied at 100 % of the recommended NPK were 407.92, 384.58 and 306.17 ppm at tillering, heading and maturity, respectively, while the respective lowest available K values of 258.50, 261.83 and 204.50 ppm were recorded for the control pots. However, there were no significant difference in available K between 66.67 % and 100 % NPK at tillering; and between 4 and 6 t VC ha⁻¹ at tillering and maturity stages of wheat. The application of vermicompost

at 2, 4 and 6 t ha⁻¹ increased availability of potassium by 15%, 30% and 34% over the control, respectively, at tillering stage. However, declines in available K were recorded at heading and maturity stages of wheat. This decline in available potassium was in exception for pots treated with 6 t ha⁻¹, which continue mineralization of K up to heading and then declined at maturity (Fig. 7). This indicates that application of vermicompost at higher rates (6 t ha⁻¹) maintains relatively high amount of available potassium up to heading stage. In line with this, Nathiya and Sanjivkumar (2015) have also found higher exchangeable K level at flowering stage than at reproductive and post harvest stages of groundnut.




Figure 5: Potassium release pattern in the soil at different growth stages of wheat as influenced by application of vermicompost




Figure 6: Potassium release pattern in the soil at different growth stages of wheat as influenced by application of chemical fertilizer

Moreover, application of potassium from chemical fertilizer had also contributed

available K to the soil solution. Applications of graded NPK at 33.33 %, 66.67 % and 100 %

of the recommended doses showed 28%, 50% and 58% increments, respectively, in available potassium contents, over the control, at tillering stage. However, the increase in available potassium due to the application of graded levels of vermicompost was reached its maximum at tillering stage, which declined slightly at heading and sharply at maturity stages (Fig. 8). This reduction at heading stage may possibly be due to the uptake by straw and grain of wheat after tillering stage. Unlike vermicompost, there were no additional organic matter on chemically fertilized pots, which let wheat to deprive only from the added nutrients and that might resulted to the excessive decrease in available forms of potassium as the crop growth excels. Likewise, K was also correlated significantly with the availability of total organic matter (Angelova *et al.*, 2013).

CONCLUSIONS AND RECOMMENDATIONS

In general, application of inorganic fertilizers as well as vermicompost at low levels have a limited capacity in supplying NPK, which might not line up with the trend of wheat nutrient uptake. As there was no addition of Nitrogen sources after the basal application, a reduction of its total amount from all treatments were evidenced throughout the growing season. However, unlike chemical fertilizers, the higher rates of vermicompost showed a continues mineralization trend on available P and K up to heading stages of wheat, which was probably due to the activity of microorganisms along with the presence of organic matter.

Consequently, Vermicompost at a rate of 6 t ha⁻¹ seems to have an equivalent release of nutrients with the recommended doses of NPK from inorganic sources. Out of all growth stages, the availability of nutrients (NPK) at tillering and heading seems to have a strong positive correlation with the uptake and yields of wheat. Hence, any soil fertility management practices that can augment the availability of

nutrients after these growth stages might not have a significant contribution to the uptake as well as to the yields of wheat.

REFERENCES

- Angelova, V R, Akova, V I, Artinova, N S. & Ivanov, K I. (2013). The effect of organic amendments on soil chemical characteristics. *Bulgarian Journal of Agricultural Science*, 19(5), 958–971.
- Arancon, N.Q., Edwards, C.A., & Bierman, P. (2006). Influences of vermicomposts on field strawberries: Part 2. Effects on soil microbiological and chemical properties. *Bioresource Technology; Elsevier*, 97, 831–840.
- ATA. (2014). *Soil Fertility Status and Fertilizer Recommendation Atlas for Tigray Regional State, Ethiopia*. MOA; ATA: Addis Abeba.
- Berhanu D. (1980). *The physical criteria and their rating proposed for land evaluation in the highland region of Ethiopia*. Land Use Planning and Regulatory Department. Ministry of Agriculture: Addis Ababa.
- Bouyoucos, G.J. (1962). Hydrometer method improved for making particle size analyses of soil. *Agronomy Journal*, 53, 464-465.
- Bremner, J.M., & Mulvaney, C.S. (1982). Nitrogen- Total. In A.L. Page, R.H. Miller & D.R. Keeney (Eds.). *Methods of soil analysis. Part 2-Chemical and microbiological properties*. *Agronomy*, 9 (2), 595-624.
- Chapman, H.D. (1965). Cation-exchange capacity. In: C. A. Black (ed.) *Methods of soil analysis - Chemical and microbiological properties*. *Agronomy*, 9, 891-901.
- Cottenie, A. (1980). Soil and plant testing as abasis of fertilizer recommendations. *FAO soil bulletin* 38/2. Food and

- Agriculture Organization of the United Nations: Rome.
- FAO. (2006). Plant nutrition for food security: A guide for integrated nutrient management. *FAO, Fertilizer and Plant Nutrition Bulletin 16*, Rome.
- Gete Z., Gete G., Agegnehu D., A. & Shahidur, R. (2010). *Fertilizer and Soil Fertility Potential in Ethiopia: Constraints and opportunities for enhancing the system*. International Food Policy Research Institute (IFPRI).
- Gupta, P.K. (2000). *Soil, Plant, Water and Fertilizer Analysis*. India.: AGROBIOS Publisher.
- Hammermeister, A M., Astatkie, T., Jeliaskova, E A., Warman, P R., & Martin, R C. (2006). Nutrient supply from organic amendments applied to unvegetated soil, lettuce and orchardgrass. *Canadian Journal of Soil Science*, 86, 21–33.
- Hazelton, P. & Murphy, B. (2007). *Interpreting soil test results: what do all the numbers mean?*, Collingwood VIC 3066, Australia: CSIRO Publishing.
- Jones, C A., Koenig, R T., Ellsworth, J W., Brown, B D. & Jackson, G D. (2007). *Management of Urea Fertilizer to Minimize Volatilization*, Montana State University, Montana.
- Lu, C., Ma, J., Chen, X., Zhang, X., Shi, Y. & Huang, B. (2010). Effect of nitrogen fertilizer and maize straw incorporation on NH_4^+ -15N and NO_3^- -15N accumulation in black soil of Northeast China among three consecutive cropping cycles. *Journal of Soil Science and Plant Nutrition*, 10(4), 443–453.
- Lungmuana, Ghosh, M. & Patra, P.K. (2013). Effect of integrated nutrient management on available phosphorus influencing grain and straw yield of rice (cv . IR-36) in an Alfisol. *Journal of Crop and Weed*, 9(1), 89–93.
- Malik, M A., Khan, K S., Marschner, P. & Fayyaz-ul-Hassan. (2013). Microbial biomass, nutrient availability and nutrient uptake by wheat in two soils with organic amendments. *Journal of Soil Science and Plant Nutrition*, 13(4), 955–966.
- Mehta, K.M., Puntamkar, S.S. & Kalamkar, V.G. (1963). Study on uptake of nutrients by wheat as influenced by nitrogen and phosphorus fertilization. *Journal of Soil Science and Plant Nutrition*, 9(5), 29–34.
- Mitiku, H., Tigist, A., Asmelash, B. & Zelalem, H. (2007). *Pedological Map Development for selected Woredas of Tigray Region*. Mekelle University, Ethiopia.
- Murphy, H.F. (1968). A report on fertility status and other data on some soils of Ethiopia. Collage of Agriculture HSIU. *Experimental Station Bulletin*, 44, 551p. Collage of Agriculture.
- Najafi-ghiri, M. (2014). Effects of Zeolite and Vermicompost Applications on Potassium Release from Calcareous Soils. *Journal of Soil and Water Research*, 9(1), 31–37.
- Nathiya, K. V., & Sanjivkumar, V. (2015). Dynamics of potassium by the combined use of organic manures and inorganic potassium fertilizers on available nutrients of groundnut crop (*Arachis hypogea*) in Madukkur soil series. *Journal of Applied and natural Science*, 7(1), 197–202.
- Olsen, S.R., Cole, C.V., Watanabe, F.S., & Dean, L.A. (1954). *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*. USDA Circular, 939.
- Rhoades, J.D. (1982). In *Methods of Soil Analysis, Part 2. Second Edition* (A.L. Page, Miller & D.R. Keeney, Eds.). *American Society of Agronomy*.

- Madison, USA.
- Roy, R N., Finck, A., Blair, G J. & Tandon, H L S. (2006). *Plant nutrition for food security: A guide for integrated nutrient management*. Rome: FAO Fertilizer and Plant Nutrition.
- Sahlemedhin S. & Taye B. (2000). *Procedures for Soil and Plant Analysis*. National Soil Research center. Ethiopian Agricultural Research Organization: Addis Ababa.
- Sheoran, H S., Duhan, B S., Grewal, K S. & Sheoran, S. (2015). Grain yield and NPK uptake of wheat (*Triticum aestivum* L.) as influenced by nitrogen, vermicompost and herbicide (Clodinafop propargyl). *African Journal of Agricultural Research*, 10(42), 3952–3961.
- Sinha, R K., Heart, S., Valani, D. & Chauhan, K. (2009). “Vermiculture and Sustainable Agriculture”. *American-Eurasian Journal of Agriculture and Environmental Science*, 5(S), .01–55.
- Tekalign T, (1991). Soil, plant, water, fertilizer, animal manure and compost analysis. *Working Document 13*. International Livestock Research Center for Africa, Addis Ababa.
- Thakare, R. & Wake, A. (2015). Dynamics of Soil Nutrients, Uptake and Yield under Organically Growth Rainfed pearl Millet in vertisol. *International Journal of Tropical Agriculture*, 33(1), 11–14.
- Tharmaraj, K., Ganesh, P., Kolanjinathan, K., Kumar, R S. & Anandan, A. (2010). Influence of Vermicompost and Vermiwash on Physio Chemical Properties of Black Gram Cultivated Soil. *International Journal of Recent Scientific Research*, 3, 77–83.
- Tirol-Padre, A., Ladha, J K., Regmi, A P., Bhandari, A L. & Inubushi, K. (2007). Organic Amendments Affect Soil Parameters in Two Long-Term Rice-Wheat Experiments. *Soil Science Society of America*, 71(2), 442–452.
- Walkley, A. & Black, I.A. (1934). An examination of the Degtjareff method for determining soil organic matter and proposed chromic acid titration method. *Soil Science*, 37, 29-38.